



# Cadmium References Data

|                         |  |
|-------------------------|--|
| <b>Title</b>            | Bioremediation of cadmium-contaminated paddy soil using an autotrophic and heterotrophic mixture   |
| <b>Author Name</b>      | Menglong Xu, Yazhi Liu, Yan Deng, Siyuan Zhang, Xiaodong Hao, Ping Zhu, Jieyi Zhou, Huaqun Yin, Yili Liang, Hongwei Liu, Xueduan Liu, Lianyang Bai, Luhua Jiang and Huidan Jiang   |
| <b>Journal Name</b>     | RSC Advances   |
| <b>Year</b>             | 2020   |
| <b>Volume and Issue</b> | 10(44)   |
| <b>Pages</b>            | 26090-26101  |
| <b>Abstracts</b>        | <p>Cadmium (Cd) pollution poses a serious risk to human health and ecological security. Bioremediation can be a promising and effective remediation technology for treating Cd contaminated soils. In this study, seven heterotrophic strains were isolated from Cd contaminated soil and 7 autotrophic strains were isolated from acid mine drainage. Cd removal efficiencies were compared after leaching with autotrophic bacteria (Att-sys), heterotrophic isolates (Htt-sys) and cooperative leaching systems (Co-sys) in laboratory agitating reactors. The results indicated that Cd removal efficiency of Co-sys (32.09%) was significantly higher than that of Att-sys (23.24%) and Htt-sys (0.74%). By analyzing the soil microbial community in different bioleaching systems, we found that the addition of heterotrophic isolates significantly promoted the growth of some heavy metal resistant inhabitants (<i>Massilia</i>, <i>Alicyclobacillus</i>, <i>Micromonospora</i>, etc.), and Co-sys had a minor effect on the growth of soil indigenous microbes. In Co-sys, the content of the four Cd fractions all decreased compared with other leaching systems. The analysis of soil physicochemical parameters during the leaching process showed that pH and ORP (oxidation reduction potential) were not the only determinants for Cd removal efficiency in Co-sys, synergistic metabolic activities of autotrophic and heterotrophic strains may be other determinants. This study demonstrated that cooperative bioremediation may prove to be a safe and efficient technique for field application in heavy metal soil pollution.</p> |
| <b>Keywords</b>         | Cadmium; Heterotrophic strains; autotrophic strains; oxidation reduction potential; bioremediation   |

|                         |   |
|-------------------------|---|
| <b>Title</b>            | <b>Toxicity of cadmium and its competition with mineral nutrients for uptake by plants: A review</b>  |
| <b>Author Name</b>      | Shiyu Qin, Hongen Liu, Zhaojun Nie, Zedren Gel, Wei Gao, Chang Li, Peng Zhao  |
| <b>Journal Name</b>     | Pedosphere  |
| <b>Year</b>             | 2020  |
| <b>Volume and Issue</b> | 30(2)   |
| <b>Pages</b>            | 168-180   |
| <b>Abstracts</b>        | <p>Cadmium (Cd) is a toxic heavy metal occurring in the environment naturally and is also generated through various anthropogenic sources and acts as a pollutant. Human health is affected by Cd pollution in farmland soils because food is the main source of Cd intake in the non-smoking population. For crops, Cd toxicity may result from a disturbance in uptake and translocation of mineral nutrients and disturbance in plant metabolism, inhibiting plant growth and development. However, plants have Cd tolerance mechanisms, including restricted Cd uptake, decreased Cd root-to-shoot translocation, enhanced antioxidant enzyme activities, and increased production of phytochelatins. Furthermore, optimal supply of mineral nutrients is one of the strategies to alleviate the damaging effects of Cd on plants and to avoid its entry into the food chain. The emerging molecular knowledge contributes to understanding Cd uptake, translocation, and remobilization in plants. In this review, Cd toxicity and tolerance mechanisms, agricultural practices to minimize Cd accumulation, Cd competition with essential elements (calcium, copper, iron, zinc, and manganese), and genes associated with Cd uptake are discussed in detail, especially regarding how these mineral nutrients and genes play a role in decreasing Cd uptake and accumulation in crop plants.</p> |
| <b>Keywords</b>         | Antagonistic interaction; Gene regulation; Heavy Metal; Mitigation measures; Tolerance; Toxicity; Uptake and Transport  |

|                         |  |
|-------------------------|--|
| <b>Title</b>            | <b>Remediation of Cadmium-Polluted Soil Using Plant Growth-Promoting Rhizobacteria and Natural Zeolite</b>   |
| <b>Author Name</b>      | V. P. Shabayev, E. A. Bocharnikova & V. E. Ostroumov   |
| <b>Journal Name</b>     | Eurasian Soil Science  |
| <b>Year</b>             | 2020   |
| <b>Volume and Issue</b> | 53   |
| <b>Pages</b>            | 809–819  |
| <b>Abstracts</b>        | <p>The impact of two strains of <i>Pseudomonas</i> bacteria and natural zeolite on the growth and elemental composition of barley plants was studied in an agrogray soil (<i>Luvisol</i>) artificially contaminated with cadmium in pot experiments. Application of <i>P. fluorescens</i> 21, or <i>P. putida</i> 23, or zeolite eliminated the heavy metal toxicity for plants. The cumulative effect of co-application of <i>P. fluorescens</i> 21 and zeolite was insignificant. The bacteria- or zeolite-mediated plant tolerance to cadmium was attributed to the enhanced root system development, decreased cadmium translocation into the roots, and improved mineral nutrition of the plants. Elevated nutrient uptake by the plants under the influence of bacteria and zeolite was the result of plant growth stimulation without significant changes in the concentrations of macronutrients N, P, K, Ca, and Mg, as well as Fe and micronutrients Zn, Mn, and Cu in plant tissues, including grain. Application of <i>P. fluorescens</i> 21 enhanced the Cd fixation in the soil organic matter in the first half of the growing season, which could be due to the metal sequestration by bacterial siderophores. Thus, application of the bacteria and natural zeolite can be recommended in the strategies for Cd-polluted soil remediation based on environment-friendly technologies.</p> |
| <b>Keywords</b>         | Cadmium Stress; Barley; Pseudomonas Bacteria; Silicon-Rich Mineral   |

|                         |   |
|-------------------------|---|
| <b>Title</b>            | <b>Responses of nitric oxide and hydrogen sulfide in regulating oxidative defence system in wheat plants grown under cadmium stress</b>   |
| <b>Author Name</b>      | Cengiz Kaya, Muhammad Ashraf, Mohammed Nasser Alyemeni and Parvaiz Ahmad  |
| <b>Journal Name</b>     | Physiologia Plantarum   |
| <b>Year</b>             | 2020  |
| <b>Volume and Issue</b> | 168   |
| <b>Pages</b>            | 345-360   |
| <b>Abstracts</b>        | <p>We conducted a study to evaluate the interactive effect of NO and H<sub>2</sub>S on the cadmium (Cd) tolerance of wheat. Cadmium stress considerably reduced total dry weight, chlorophyll a and b content and ratio of Fv/Fm by 36.7, 48.6, 26.7 and 19.5%, respectively, but significantly enhanced the levels of hydrogen peroxide (H<sub>2</sub>O<sub>2</sub>) and malondialdehyde (MDA), endogenous H<sub>2</sub>S and NO, and the activities of antioxidant enzymes. Exogenously applied sodium nitroprusside (SNP) and sodium hydrosulfide (NaHS), donors of NO and H<sub>2</sub>S, respectively, enhanced total plant dry matter by 47.8 and 39.1%, chlorophyll a by 92.3 and 61.5%, chlorophyll b content by 29.1 and 27.2%, Fv/Fm ratio by 19.7 and 15.2%, respectively, and the activities of antioxidant enzymes, but lowered oxidative stress and proline content in Cd-stressed wheat plants. NaHS and SNP also considerably limited both the uptake and translocation of Cd, thereby improving the levels of some key mineral nutrients in the plants. Enhanced levels of NO and H<sub>2</sub>S induced by NaHS were reversed by hypotuarine application, but they were substantially reduced almost to 50% by cPTIO (a NO scavenger) application. Hypotuarine was not effective, but cPTIO was highly effective in reducing the levels of NO and H<sub>2</sub>S produced by SNP in the roots of Cd-stressed plants. The results showed that interactive effect of NO and H<sub>2</sub>S can considerably improve plant resistance against Cd toxicity by reducing oxidative stress and uptake of Cd in plants as well as by enhancing antioxidative defence system and uptake of some essential mineral nutrients.</p> |
| <b>Keywords</b>         | Hydrogen Peroxide; Malondialdehyde; Sodium Nitroprusside; Sodium Hydrosulfide; Chlorophyll; Oxidative Stress  |

|                         |   |
|-------------------------|---|
| <b>Title</b>            | <b>Mechanisms of Cadmium Accumulation in Plants</b>   |
| <b>Author Name</b>      | Thibault Sterckeman & Sébastien Thomine   |
| <b>Journal Name</b>     | Critical Reviews in Plant Sciences  |
| <b>Year</b>             | 2020  |
| <b>Volume and Issue</b> | 39(4)   |
| <b>Pages</b>            | 322-359   |
| <b>Abstracts</b>        | <p>Cadmium is a non-essential trace metal, which is highly toxic to nearly all living organisms. Soil pollution causes Cd contamination of crops, thereby rendering plant products responsible for the chronic low level Cd over-exposure of numerous populations in the world. For this reason, Cd accumulation in plants has been studied for about five decades now. The research first focused on the relationships between plant and soil Cd levels, on the factors of the metal availability in soil, as well as the root uptake processes. Cd distribution in plant organs was also investigated, first using a macroscopic and eco-physiological approach, and then with the help of molecular biology tools, at both tissue and cell scales. Cadmium has no biological function and hijacks the transport pathways of micronutrients such as Fe, Mn, or Zn, in order to enter the plant through the roots and be distributed to all its organs. The study of the genes that control the influx and efflux of the Cd<sup>2+</sup> ion in the cytosol, vacuoles, and vascular tissues has significantly contributed to the understanding of the metal root uptake and of its transfer to the aerial parts. However, the mechanisms responsible for its distribution to the different above-ground tissues and specially to fruits and seeds have yet to be clarified. This review summarizes current knowledge in order to present a detailed overview of Cd transport and storage, from the rhizosphere to the different organs and tissues of the plant.</p> |
| <b>Keywords</b>         | Cell wall; chelation; inter-element competition; ion transporter; iron; manganese; nitrogen; organic acid; phloem; phytochelatin; silicon; speciation; vacuolar sequestration; xylem sap; zinc  |

|                         |   |
|-------------------------|---|
| <b>Title</b>            | Potential use of king grass ( <i>Pennisetum purpureum</i> Schumach. × <i>Pennisetum glaucum</i> (L.) R.Br.) for phytoextraction of cadmium from fields  |
| <b>Author Name</b>      | Zhiqiang Zhou, Yangyang Guo, Li Hu, Lan He, Bo Xu, Zhenrui Huang, Guo Wang & Yanhui Chen  |
| <b>Journal Name</b>     | Environmental Science and Pollution Research  |
| <b>Year</b>             | 2020  |
| <b>Volume and Issue</b> | 27  |
| <b>Pages</b>            | 35249–35260   |
| <b>Abstracts</b>        | <p>Using king grass (<i>Pennisetum purpureum</i> Schumach. × <i>Pennisetum glaucum</i> (L.) R.Br.) for phytoextraction is a promising technology for producing large amounts of biomass fuel while remediating contaminated soil. To assess the practical phytoextraction capacity of king grass, we conducted a field experiment with three different soil types (loam, sandy loam, clay loam) and cadmium (Cd) concentrations (0, 0.25, 0.5, 1, 2, 4, 8, and 16 mg kg<sup>-1</sup>, aged stably for 6 years). King grass were harvested at two different periods (elongation and maturity) to identify the optimal harvest time for extraction efficiency. The results showed that all treatments had bioconcentration factor (BCF) &gt; 1 and translocation factor (TF) &lt; 1; Cd is mainly stored in the roots. However, due to a high shoot biomass, the highest quantity of Cd extracted from shoots was 2.75 mg plant<sup>-1</sup>, from the experimental group with 16 mg kg<sup>-1</sup> Cd added in sandy loam. A significant positive relationship (P &lt; 0.05) was observed between the amount of Cd extracted from king grass stems, leaves, and roots from soil with the diethylene triamine pentacetate acid (DTPA) extractable Cd concentration. The Cd concentration in shoots at the maturity stage is lower than at the elongation stage, mainly due to the effect of biological dilution. Meanwhile, there is significantly more biomass (P &lt; 0.05) at the maturity stage than at the elongation stage. At the latter, the extraction efficiency of the three soils was loam &gt; sandy loam &gt; clay loam, while at maturity it was sandy loam &gt; clay loam &gt; loam. This change in extraction efficiency can be attributed mainly to differences in soil DTPA-extractable Cd concentration and growth rate caused by differences in soil physical and chemical properties. According to calculations from multiple harvests using three types of soil, remediating contaminated soil with 0–16 mg kg<sup>-1</sup> Cd would take 13.9–224.5 and 19.5–250.6 years, extracting 7.21–265.23 and 4.96–330.52 g ha<sup>-1</sup> Cd while producing 33.62–66.50 and 73.8–110.5 t ha<sup>-1</sup> dry biomass at the elongation (90 days) and maturity (120 days) stages, respectively. In summary, king grass has major potential for remediating Cd-contaminated soil while producing large volumes of biofuel.</p> |
| <b>Keywords</b>         | Cadmium; Growth period; Phytoextraction; <i>Pennisetum purpureum</i> Schumach. × <i>Pennisetum glaucum</i> (L.) R.Br.; Soil type  |

|                         |   |
|-------------------------|---|
| <b>Title</b>            | Effects of intercropping accumulator plants and applying their straw on the growth and cadmium accumulation of <i>Brassica chinensis</i> L.   |
| <b>Author Name</b>      | Yi Tang, Liming Wang, Yongdong Xie, Xuena Yu, LiJin Lin, Huanxiu Li, Ming'an Liao, Zhihui Wang, Guochao Sun, Dong Liang, Hui Xia, Xun Wang & Lihua Tu   |
| <b>Journal Name</b>     | Environmental Science and Pollution Research  |
| <b>Year</b>             | 2020  |
| <b>Volume and Issue</b> | 27  |
| <b>Pages</b>            | 39094–39104   |
| <b>Abstracts</b>        | <p>Two pot experiments were conducted to study the effects of intercropping cadmium (Cd) accumulator plants (<i>Stellaria media</i> (L.) Villars, Cardamine hirsuta, Cerastium glomeratum Thuill, and <i>Galium aparine</i> L.) and applying their straw on the growth and Cd accumulation of <i>Brassica chinensis</i> L. Intercropping with four accumulator plants reduced the biomass, water content, and photosynthetic pigment content of <i>B. chinensis</i> compared with monoculture. Intercropping with accumulator plants increased the Cd content in the roots and shoot of <i>B. chinensis</i>, and the translocation factor (TF), root bioconcentration factor (root BCF), and shoot bioconcentration factor (Shoot BCF) increased. The soil pH decreased and the soil available Cd content increased by intercropping. Thus, intercropping with four accumulator plants can promote the Cd uptake of <i>B. chinensis</i>. The straw of four accumulator plants reduced the biomass, water content, and photosynthetic pigment content of <i>B. chinensis</i> compared with the control. The straw of <i>S. media</i> and <i>C. hirsute</i> increased the Cd content in the roots and shoots of <i>B. chinensis</i>, TF, root BCF, and shoot BCF. The straw of <i>C. glomeratum</i> and <i>G. aparine</i> decreased the Cd content in the roots and shoots of <i>B. chinensis</i>, TF, root BCF, and shoot BCF. The soil pH increased and the soil available Cd content decreased by application of straw. Thus, the straw of <i>C. glomeratum</i> and <i>G. aparine</i> can reduce the Cd uptake of <i>B. chinensis</i>.</p> |
| <b>Keywords</b>         | Accumulator plants; Intercropping; Straw; <i>Brassica chinensis</i> L; Cadmium  |

|                         |  |
|-------------------------|--|
| <b>Title</b>            | <b>Cadmium Stress and Toxicity in Plants: An Overview</b>  |
| <b>Author Name</b>      | Bala Murugan Shanmugaraj, AshwiniMalla & Sathishkumar Ramalingam   |
| <b>Journal Name</b>     | Cadmium Toxicity and Tolerance in Plants (From Physiology to Remediation)  |
| <b>Year</b>             | 2019   |
| <b>Volume and Issue</b> | 1  |
| <b>Pages</b>            | 1 - 17   |
| <b>Abstracts</b>        | Heavy metal pollution has long been a major environmental problem, and threatens all living forms globally. Cadmium (Cd) is one of the nonessential, highly toxic environmental pollutants worldwide that causes deleterious effects and serious problems in agriculture. Plants growing in Cd-contaminated soil uptake the heavy metal through their roots, which accumulates in different organs, eventually reducing plant growth and productivity. Bioaccumulation of high concentrations of Cd in plants enters the food chain and affects both animals and humans. In plants, several metabolic processes are associated with Cd toxicity and its tolerance. This chapter summarizes the effects of Cd toxicity in plant growth and other related physiological and metabolic processes, including an overview of phytoremediation technologies to clean up Cd contamination in the environment. |
| <b>Keywords</b>         | Abiotic stress; Accumulation; Cadmium; Heavy metals; Phytoremediation; Toxicity  |



|                         |  |
|-------------------------|--|
| <b>Title</b>            | <b>Characteristics of cadmium accumulation and isotope fractionation in higher plants</b>  |
| <b>Author Name</b>      | Rongfei Wei, Qingjun Guo, Liyan Tian, Jing Kong, Yang Baia,Chukwunonso Peter Okoli & Liyuan Wang   |
| <b>Journal Name</b>     | Ecotoxicology and Environmental Safety   |
| <b>Year</b>             | 2019   |
| <b>Volume and Issue</b> | Volume 174   |
| <b>Pages</b>            | 1 -11  |
| <b>Abstracts</b>        | Cadmium (Cd) pollution of the soil is an important global environmental issueowing to its great toxicity. The study of metal isotope fractionation is a novel technique that could be used to identify and quantify metal uptake and transport mechanisms in plant. In this study, cadmium tolerant <i>Ricinus communis</i> and hyperaccumulator <i>Solanum nigrum</i> have been cultured in different Cd concentration nutrient solutions. The Cd isotope values, metal elements concentrations in the organs (root, stem and leaf) in the two plant species have been measured during the growth periods (10d, 15d, 20d, 25d, and 30d). The results indicate that the organs of <i>S. nigrum</i> could be enriched with lighter Cd isotopes compared with <i>R. communis</i> . In addition, the Cd isotope fractionation become smaller when the plants were subjected to high Cd toxicity, which indicates that Cd isotope fractionation reflected the extent of Cd toxicity to plants. This study advances our current view of Cd translocation machination in plants. |
| <b>Keywords</b>         | Cd; Isotope fractionation; <i>Ricinus communis</i> ; <i>Solanum nigrum</i> ; Plants  |

|                         |   |
|-------------------------|---|
| <b>Title</b>            | <b>Phytoremediation of cadmium-polluted soils with <i>Ipomoea asarifolia</i> (Desr.) Roem. &amp; Schult</b>   |
| <b>Author Name</b>      | S Shehu, R.S.U. Wasagu, S.A. Anka, J.C. Okoro & Y Saidu   |
| <b>Journal Name</b>     | Journal of Applied Sciences and Environmental Management  |
| <b>Year</b>             | 2019  |
| <b>Volume and Issue</b> | 23, 2   |
| <b>Pages</b>            | 253 - 259   |
| <b>Abstracts</b>        | <p>Phytoremediation is an alternative method for restoring soils polluted with heavy metals which is cost-effective and environment-friendly. The present study evaluated the potential of <i>Ipomoea asarifolia</i> to remediate soils experimentally-amended with Cadmium. The plant was grown on soils amended with 0, 1500, 2000, and 2500 mg CdCl<sub>2</sub> salt. The salt was mixed with small portions of the soils and made upto 3kg salt/soil mixtures each. These were applied into 4 separate polythene-pots labelled; A, B, C and D respectively. Sample A containing 3kg non-amended soil (without Cd) served as the control. The concentrations of Cd applied to the soils were therefore; 0, 306.61, 408.82 and 511.02 mg/kg soils in the samples A-D respectively. Atomic absorption spectrophotometry (AAS) was used to analyse the bioaccumulation of Cd in the plant's parts, over three harvesting phases of the study period. The results revealed that <i>I. asarifolia</i> is a good phytoaccumulator as it accumulated a total biomass of <math>0.23 \pm 0.63</math>, <math>272.85 \pm 1.99</math>, <math>377.40 \pm 0.63</math> and <math>459.48 \pm 0.60</math> mg/kg Cd from the amended soils A-D respectively. The Transportation Indices; RTI and STI for translocation of Cd to the plant's stems and leaves were both greater than 1 (TI &gt;1), indicating that the plant has a phytoextraction potential for Cadmium. These results therefore, suggest that <i>I. asarifolia</i> could be effective in phytoremediation of Cadmium-polluted environments.</p> |
| <b>Keywords</b>         | Heavy metals; cadmium; pollution; phytoremediation; <i>Ipomoea asarifolia</i>   |

|                         |  |
|-------------------------|--|
| <b>Title</b>            | <b>Biochar facilitated the phytoremediation of cadmium contaminated sediments: Metal behavior, plant toxicity, and microbial activity</b>  |
| <b>Author Name</b>      | Xiaomin Gong, Danlian Huang, Yunguo Liu, Guangming Zeng, Sha Chen, Rongzhong Wang, Piao Xu, Min Cheng, Chen Zhang & Wenjing Xue  |
| <b>Journal Name</b>     | Science of The Total Environment   |
| <b>Year</b>             | 2018   |
| <b>Volume and Issue</b> | Volume 666   |
| <b>Pages</b>            | 1126-1133  |
| <b>Abstracts</b>        | Cadmium (Cd) contamination in river sediments becomes increasingly serious, and phytoremediation has been used to remediate Cd contaminated sediments, but the remediation efficiency needs to be improved. In this study, tea waste derived biochar (TB) was used to facilitate the phytoremediation of Cd contaminated sediments. Results showed that TB at 100, 500 and 1000 mg kg <sup>-1</sup> increased Cd accumulation and translocation in ramie seedlings by changing Cd speciation in sediments and altering the subcellular distribution of Cd in plant cells. TB at low contents alleviated Cd induced toxicity in ramie seedlings by promoting plant growth and mitigating the oxidative stress. In addition, the activities of urease-, phosphatase-, and catalase-producing microbes in the Cd contaminated sediments were promoted by the application of TB. These findings demonstrated that biochar at low concentrations could improve the phytoremediation efficiency and mitigating Cd-induced toxicity to plants and microbes in Cd contaminated sediments. This study herein provides a novel technological application of waste biomass in controlling and mitigating risks of heavy metals. |
| <b>Keywords</b>         | Phytoremediation; Cadmium; Plants; Microbes; Sediments   |

|                         |   |
|-------------------------|---|
| <b>Title</b>            | <b>Sulfide alleviates cadmium toxicity in Arabidopsis plants by altering the chemical form and the subcellular distribution of cadmium</b>  |
| <b>Author Name</b>      | Mei Yan,Guan,Hai Hua Zhang,Wei Pan,Chong Wei Jin, Xian Yong Lin   |
| <b>Journal Name</b>     | Science of The Total Environment  |
| <b>Year</b>             | 2018  |
| <b>Volume and Issue</b> | Volume 627  |
| <b>Pages</b>            | 663-670   |
| <b>Abstracts</b>        | Several sulfur compounds are thought to play important roles in the plant tolerance to cadmium (Cd), but the role of inorganic sulfide in Cd tolerance remains largely unknown. In this study, we found that Cd exposure increased the accumulation of soluble sulfide in Arabidopsis plants. When exogenous sulfide, in the form of NaHS, was foliarly applied, Cd-induced growth inhibition and oxidative stress were alleviated. In addition, although the foliar application of sulfide did not affect the total Cd levels, it significantly decreased the soluble Cd fractions in plants. Furthermore, foliar applications of sulfide decreased Cd distribution in the cytoplasm and organelles, but increased Cd retention in the cell wall, which is a less sensitive compartment. These results suggest that the Cd-induced accumulation of soluble sulfide alleviates Cd toxicity in plants by inactivating Cd and sequestering it into the cell wall. |
| <b>Keywords</b>         | Arabidopsis thaliana, Cadmium, Sulfide, Cadmium tolerance   |

|                         |  |
|-------------------------|--|
| <b>Title</b>            | Selenium mitigates cadmium-induced oxidative stress in tomato ( <i>Solanum lycopersicum</i> L.) plants by modulating chlorophyll fluorescence, osmolyte accumulation, and antioxidant system.  |
| <b>Author Name</b>      | Mohammed Nasser Alyemeni, Mohammad Abass Ahanger, Leonard Wijaya, Pravej Alam, Renu Bhardwaj, Parvaiz Ahmad  |
| <b>Journal Name</b>     | Protoplasma  |
| <b>Year</b>             | 2017   |
| <b>Volume and Issue</b> | --   |
| <b>Pages</b>            | 1-11   |
| <b>Abstracts</b>        | <p>Pot experiments were conducted to investigate the role of selenium in alleviating cadmium stress in <i>Solanum lycopersicum</i> seedlings. Cadmium (<math>150 \text{ mg L}^{-1}</math>) treatment caused a significant reduction in growth in terms of height and biomass accumulation and affected chlorophyll pigments, gas exchange parameters, and chlorophyll fluorescence. Selenium (<math>10 \text{ }\mu\text{M}</math>) application mitigated the adverse effects of cadmium on growth, chlorophyll and carotenoid contents, leaf relative water content, and other physiological attributes. Lipid peroxidation and electrolyte leakage increased because of cadmium treatment and selenium-treated plants exhibited considerable reduction because of the decreased production of hydrogen peroxide in them. Cadmium-treated plants exhibited enhanced activity of antioxidant enzymes that protected cellular structures by neutralizing reactive free radicals. Supplementation of selenium to cadmium-treated plants (Cd+Se) further enhanced the activity of superoxide dismutase (SOD), catalase (CAT), ascorbate peroxidase (APX), and glutathione reductase (GR) by 19.69, 31.68, 33.14, and 54.47%, respectively. Osmolytes, including proline and glycine betaine, increased with selenium application, illustrating their role in improving the osmotic stability of <i>S. lycopersicum</i> under cadmium stress. More importantly, selenium application significantly reduced cadmium uptake. From these results, it is clear that application of selenium alleviates the negative effects of cadmium stress in <i>S. lycopersicum</i> through the modifications of osmolytes and antioxidant enzymes.</p> |
| <b>Keywords</b>         | <i>Solanum lycopersicum</i> , Cadmium , Growth , Chlorophyll fluorescence , Proline , Lipid peroxidation , Antioxidants , Selenium   |

|                         |  |
|-------------------------|--|
| <b>Title</b>            | <b>Uptake, sequestration and tolerance of Cadmium at cellular levels in the hyperaccumulator plant species <i>Sedum alfredii</i></b>   |
| <b>Author Name</b>      | Shengke Tian Ruohan Xie Haixin Wang Yan Hu Dandi Hou Xingcheng LiaoPatrick H. Brown Hongxia Yang Xianyong Lin John M. Labavitch  |
| <b>Journal Name</b>     | Journal of Experimental Botany   |
| <b>Year</b>             | 2017   |
| <b>Volume and Issue</b> | 68 Issue 9   |
| <b>Pages</b>            | 2387-2398  |
| <b>Abstracts</b>        | <i>Sedum alfredii</i> is one of a few plant species known to hyperaccumulate cadmium (Cd). Uptake, localization, and tolerance of Cd at cellular levels in shoots were compared in hyperaccumulating (HE) and non-hyperaccumulating (NHE) ecotypes of <i>Sedum alfredii</i> . X-ray fluorescence images of Cd in stems and leaves showed only a slight Cd signal restricted within vascular bundles in the NHEs, while enhanced localization of Cd, with significant tissue- and age-dependent variations, was detected in HEs. In contrast to the vascular-enriched Cd in young stems, parenchyma cells in leaf mesophyll, stem pith and cortex tissues served as terminal storage sites for Cd sequestration in HEs. Kinetics of Cd transport into individual leaf protoplasts of the two ecotypes showed little difference in Cd accumulation. However, far more efficient storage of Cd in vacuoles was apparent in HEs. Subsequent analysis of cell viability and hydrogen peroxide levels suggested that HE protoplasts exhibited higher resistance to Cd than those of NHE protoplasts. These results suggest that efficient sequestration into vacuoles, as opposed to rapid transport into parenchyma cells, is a pivotal process in Cd accumulation and homeostasis in shoots of HE <i>S. alfredii</i> . This is in addition to its efficient root-to-shoot translocation of Cd. |
| <b>Keywords</b>         | Cadmium, Fluorescence microscopy, localized, micro X ray fluorescence, protoplasts, tolerance, Vacuole   |

|                         |  |
|-------------------------|--|
| <b>Title</b>            | Salicylic acid-induced protection against cadmium toxicity in wheat plants   |
| <b>Author Name</b>      | F.M.Shakirova,Ch.R.Allagulova,D.R.Maslennikova,E.O.Klyuchnikova,A.M.Avalbaev,M.V.Bezrukova   |
| <b>Journal Name</b>     | Environmental and Experimental Botany  |
| <b>Year</b>             | 2016   |
| <b>Volume and Issue</b> | Volume 122   |
| <b>Pages</b>            | 19-28  |
| <b>Abstracts</b>        | <p>We have studied the influence of pretreatment of wheat seedlings (<i>Triticum aestivum</i> L.) with 50 <math>\mu</math>M salicylic acid (SA) on plant resistance to subsequent action of 1 mM cadmium acetate. SA pretreatment decreased the extent of detrimental effect of cadmium on wheat plants, as judged by the decline in the level of stress-induced accumulation of MDA and electrolyte leakage. Furthermore, SA-pretreatment contributed to maintenance of growth characteristics of wheat seedlings at the level close to the control under stress conditions and to acceleration of growth recovery during post-stress period. Detected defense effect of SA may be due to a decline in the amplitude of cadmium-induced accumulation of abscisic acid (ABA) and to reduced fall of indoleacetic acid (IAA) and cytokinins (CK) in stressed plants. In the course of one day treatment, SA activated phenylalanine ammonia-lyase (PAL), the key enzyme of lignin biosynthesis, in roots of seedlings under normal growth conditions contributing to the strengthening of carrier functions of cell walls. This assumption is supported by the data showing significant decline in cadmium accumulation in SA-pretreated. Detected defense effect of SA may be due to a decline in the amplitude of cadmium-induced accumulation of abscisic acid (ABA) and to reduced fall of indoleacetic acid (IAA) and cytokinins (CK) in stressed plants. In the course of one day treatment, SA activated phenylalanine ammonia-lyase (PAL), the key enzyme of lignin biosynthesis, in roots of seedlings under normal growth conditions contributing to the strengthening of carrier functions of cell walls. This assumption is supported by the data showing significant decline in cadmium accumulation in SA-pretreated.</p> |
| <b>Keywords</b>         | Abscisic acid, Cadmium stress, Dehydrins, Hormonal balance, Phenylalanine ammonia-lyase, Lignin deposition, Plant growth, Salicylic acid<br>Wheat ( <i>Triticum aestivum</i> L.)   |

|                         |  |
|-------------------------|--|
| <b>Title</b>            | <b>Modulation and significance of nitrogen and sulfur metabolism in cadmium challenged plants</b>  |
| <b>Author Name</b>      | M. Iqbal R. Khan, Noushina Iqbal, Asim Masood, Mohammad Mobin, Naser A. Anjum  |
| <b>Journal Name</b>     | Plant Growth Regulation  |
| <b>Year</b>             | 2016   |
| <b>Volume and Issue</b> | 78, Issue1   |
| <b>Pages</b>            | 1–11   |
| <b>Abstracts</b>        | <p>As a result of rapidly increasing anthropogenic activities, input of varied metal (loids) such as cadmium (Cd) to worldwide agricultural soils and its subsequent accumulation, and obvious toxicity in plants are increasing. The role of mineral nutrients in the mitigation of Cd-accrued consequences in plants has been credibly suggested. In isolated studies, two essential mineral nutrients such as nitrogen (N) and sulfur (S) have been reported to minimize Cd-impacts in plants, and improve overall plant growth, metabolism and productivity under Cd-exposure. However, the information on the significance of N and S metabolism, and also on cross-talks on the coordination therein in Cd-challenged plants is lacking. Given the highlighted lacunae, in the light of recent research outcomes, the present review attempts to: (a) overview Cd in soil, and its major toxicity and mitigation avenues in plants, (b) appraise Cd-mediated modulation of N and S metabolism, (c) summarize the role of exogenously-sourced N and S for the mitigation of Cd toxicity, (d) critically discuss the significance of coordination between N and S metabolism for Cd-impact-mitigation, and finally to (e) highlight the major aspects to explore in the current context. The literature appraised herein suggests that a fine coordination among major pathways of N and S assimilation can enhance defense metabolites and enzymes that in turn can strengthen overall defense system, and efficiently mitigate Cd-impacts in plants. However, efforts are required to get more insights into the mechanism(s) of (co)regulation of sulfate and nitrate assimilation at the molecular level. Additionally, molecular approaches should be narrowed to enhance the production of thiols, and their products in plants through manipulating major enzymes involved in sulfate and nitrate assimilation in plants under Cd-challenged environment.</p> |
| <b>Keywords</b>         | Cadmium-phytotoxicity, Nitrogen metabolism, Sulfur metabolism, Plant-cadmium tolerance   |